

DITIONS

Cyprus  
g House, Beijing

mbH, Munich  
g, Athens

Naka-Ku Nagoya

Paulo

NUAL

ATRICS

ICAL INFORMATION—HOME EDITION

profit basis as a service to the scientific com-

SEVENTEENTH EDITION

# THE MERCK MANUAL

OF  
DIAGNOSIS AND THERAPY

*Editors*

MARK H. BEERS, M.D., and ROBERT BERKOW, M.D.

*Senior Assistant Editors*

ROBERT M. BOGIN, M.D., and  
ANDREW J. FLETCHER, M.B., B.Chir.

*Editorial Board*

Philip K. Bondy, M.D.  
Preston V. Dilts, Jr., M.D.  
Douglas A. Drossman, M.D.  
L. Jack Faling, M.D.  
Eugene P. Frenkel, M.D.  
Glen O. Gabbard, M.D.  
Robert A. Hoeckelman, M.D.  
Gerald L. Mandell, M.D.  
Fred Plum, M.D.  
G. Victor Rossi, Ph.D.  
Paul H. Tanser, M.D., F.R.C.P.(C)

Published by MERCK RESEARCH LABORATORIES

Division of Merck & Co., Inc.  
Whitehouse Station, N.J.

1999

BEST AVAILABLE COPY

# THE MERCK MANUAL

CENTENNIAL EDITION

1st Edition - 1899  
2nd Edition - 1901  
3rd Edition - 1905  
4th Edition - 1911  
5th Edition - 1923  
6th Edition - 1934  
7th Edition - 1940  
8th Edition - 1950  
9th Edition - 1956  
10th Edition - 1961  
11th Edition - 1966  
12th Edition - 1972  
13th Edition - 1977  
14th Edition - 1982  
15th Edition - 1987  
16th Edition - 1992  
17th Edition - 1999

**Editorial and Production Staff**

*Executive Editor:* Keryn A.G. Lane

*Senior Staff Editor:* Susan T. Schindler

*Staff Editor:* Julie Kostecky

*Staff Editor:* Sandra J. Masse

*Production Editor:* Debra G. Share

*Contributing Editor:* Roger I. Schreck, M.D.

*Designer:* Lorraine B. Kilmer

*Indexer:* Susan Thomas, Ph.D.

*Textbook Production Coordinator:* Diane C. Zenker

*Medical Textbook Coordinator:* Dorothy A. Bailey

*Executive Assistant:* Diane Colner-Bobrin

*Publisher:* Gary Zelko

*Advertising and Promotional Supervisor:* Pamela J. Barnes

Library of Congress Catalog Card Number 1-31760

ISBN 0911910-10-7

ISSN 0076-6526

Copyright 1999 by Merck & Co., Inc.

All rights reserved. No part of this book may be reproduced or used in any form or by any means, electronic or mechanical, including photocopying, or by any information storage and retrieval system, without permission in writing from the Publisher. Inquiries should be addressed to The Merck Manuals Department, P.O. Box 4, Merck & Co., West Point, PA 19486.

Printed in the U.S.A.

## IS DISEASES

## CHAPTER 153 - ANTIBACTERIAL DRUGS / 1123

ly in a dose of 25 to 30 mg/kg (maximum 2 g) daily in 2 divided doses as used in adults and in children.

r sulfadiazine to prevent (76). Sulfadiazine ocular

## 1- OLE

le (TMP) of the two. The dose ratio of 1:5, which is used in both. Both are active against the acid. TMP is used to treat against negative or anaerobes. usually resist-

id SMX are secreted into the milk. Both are well into the CSF. Both are active in UTI. Both are effective in treating only in children given for choice for infant pneumonia infection in adults with treatment of typhoid and d. TMP-SMX are ototoxic and ardia infec-

tion, otitis media, and acute exacerbation of chronic bronchitis.

**Adverse reactions:** The adverse reactions are the same as those listed above for sulfonamides. TMP causes identical adverse reactions to SMX but less commonly. It causes, nausea, vomiting, rash, and folate deficiency (resulting in macrocytic anemia) most likely occur. AIDS patients have a high incidence of adverse effects, especially neutropenia.

**Administration and dosage:** The oral dosage in adults is 2 regular-strength tablets (each tablet contains 80 mg TMP and 400 mg SMX) or one double-strength tablet (160 mg TMP and 800 mg SMX) bid. The usual oral dosage in children is 8 mg/kg TMP and 40 mg/kg SMX daily in 2 divided doses. The IV dosage in adults and children is 12 mg/kg TMP and 40 to 60 mg/kg SMX in 4 divided doses. Single-dose therapy with 1 or 2 double-strength tablets has been successfully used for lower UTI in women.

Much higher dosages (20 mg/kg/day TMP and 100 mg/kg/day SMX in 4 divided doses) are used in the treatment of *P. carinii* pneumonia. Much lower dosages (40 mg TMP and 200 mg SMX each night) are used for prophylaxis of UTI. For prophylaxis of *P. carinii* pneumonia, 160 mg TMP and 800 mg SMX daily or 3 days/wk (for children, 5 mg/kg/day in 2 divided doses daily or 3 times daily) is used.

### Trimethoprim

For patients allergic to sulfonamides, TMP alone has been used mainly in the treatment of chronic bacterial prostatitis and in the prophylaxis and treatment of UTI. The pharmacology and adverse effects are listed under TMP-SMX, above. The dosage in adults for treatment of UTI is 100 mg po q 12 h and 200 mg once daily.

## ANTIMICROBIAL CHEMOPROPHYLAXIS

The term antimicrobial chemoprophylaxis refers to the timely and judicious use of antimicrobial drugs to prevent infection of a symptomatic disease caused by microorganisms. Although antimicrobial drugs play a pivotal role in chemoprophylaxis, immunologic defenses also contribute to the process.

Successful prophylaxis requires either target organisms that are unlikely to develop resistance to the drugs used or clinical situations in which the duration of risk is measured in hours or days, thus permitting the effective use of antimicrobial drugs before resistance can emerge.

### Prophylaxis Against Exogenous Pathogens

Prevention of infection by exogenous pathogens, organisms that are not usually part of the normal human flora, may involve destroying microbes before attachment to host cells, modifying the microbe to prevent attachment, or eradicating colonization before either tissue invasion or toxin production begins. Indications for prophylaxis include prevention of infections caused by *Streptococcus pyogenes* in patients with a history of rheumatic fever or rheumatic heart disease (see Ch. 270) and in those with recurrent cellulitis (see Ch. 112). Patients with animal and human bites are frequently treated. The choice of drug is controversial, but amoxicillin 500 mg/clavulanate 125 mg bid for 5 days is often recommended for dog, cat, and human bites. Amoxicillin 500 mg qid or doxycycline 100 mg bid po is effective in preventing Lyme disease after tick bites; however, prophylaxis is seldom indicated because the risk of infection is generally low (see LYME DISEASE in Ch. 157).

Although *Neisseria meningitidis* and *Haemophilus influenzae* type b may occasionally colonize the upper respiratory tract of healthy persons, some persons exposed to patients with meningitis or other invasive diseases caused by these bacteria warrant prophylaxis. Household contacts, children in contact in day care centers, and others exposed to respiratory secretions of patients with invasive meningococcal disease (eg, physicians performing mouth-to-mouth resuscitation) should receive rifampin 600 mg bid for 2 days if > 12 yr of age. Pediatric dosage is 5 mg/kg q 12 h po for 2 days if < 12 yr old and 10 mg/kg q 12 h po (maximum 10 mg) for 2 days if 1 mo to 12 yr old. Ceftriaxone 125 mg IM for children < 12 yr and ceftriaxone 250 mg IM for adults are other options. Alternatively, a single dose of ciprofloxacin 500 mg po can be given to adults older than 18 yr. If unvaccinated household contacts of patients with invasive disease

caused by *H. influenzae* type b are children < 1 yr of age, all household members except pregnant women should receive oral therapy with rifampin 20 mg/kg (maximum 600 mg) daily for 4 days if > 1 mo old and 10 mg/kg for 4 days if < 1 mo old. This regimen should also be used with concurrent vaccine administration for susceptible children in day care centers, especially when two or more cases occur within 60 days and contacts include unvaccinated children < 2 yr old.

Persons traveling to endemic areas for malaria should receive prophylaxis (see Ch. 161). Although numerous studies have demonstrated that antimicrobial prophylaxis can reduce the frequency of travelers' diarrhea when traveling to high-risk areas for short durations, prophylaxis is controversial because of emerging resistance. Successful prophylaxis for this indication requires some knowledge of local susceptibility patterns for the target pathogens, usually toxin-producing strains of *Escherichia coli*. Most authorities prefer to reserve antimicrobial therapy for persons who develop diarrhea. For persons who must have prophylaxis, ciprofloxacin 500 mg/day po, norfloxacin 400 mg/day po, or two bismuth subsalicylate tablets po qid are effective. In some areas with low rates of antimicrobial resistance, oral trimethoprim-sulfamethoxazole (TMP-SMX) 160 mg/day (TMP) and 800 mg/day (SMX) or doxycycline 100 mg/day may be effective.

Antimicrobial prophylaxis is also used to prevent some viral infections. Prophylaxis with amantadine or rimantadine is effective during influenza A epidemics (see Ch. 162). Similarly, those sustaining parenteral exposures to blood or certain other body fluids from patients infected with HIV or those exposed to laboratory stocks of HIV may require prophylaxis with antiretroviral drugs (see Ch. 163).

Certain other uses of chemoprophylaxis are recommended, even though evidence from clinical trials is lacking. For the prevention of plague in exposed persons and laboratory workers, tetracycline 30 mg/kg/day po in 4 divided doses for 10 days or streptomycin 1 g IM daily for 1 wk appears to be effective. For prevention of pertussis in close contacts or colonized persons, erythromycin in an oral dose of 12.5 mg/kg qid (not to exceed 2 g/day) for 14 days is used. Chemoprophylaxis is widely used to prevent sexually transmitted diseases in the contacts of

## 1124 / SECTION 13 - INFECTIOUS DISEASES

active cases as well as in rape victims. Sexual partners of patients with gonorrhea, chancroid, lymphogranuloma venereum, nongonococcal urethritis, syphilis, trichomoniasis, scabies, pediculosis pubis, and infection caused by *Chlamydia trachomatis* receive standard therapeutic courses of the drugs normally used to treat these infections whether symptoms are present or not (see Ch. 164). Rape victims are treated with single doses of ceftriaxone 250 mg IM and metronidazole 2 g po and with doxycycline 100 mg po bid for 7 days. A single topical application of a 1% silver nitrate, 0.5% erythromycin ophthalmic ointment, or 1% solution of tetracycline ophthalmic ointment is used to prevent gonococcal and chlamydial infections in the eyes of newborns (see Ch. 260). Penicillin V, 20 mg/kg/day in 2 divided doses for children < 5 years and 250 mg po bid for children > 5 years, or amoxicillin, 125 mg po bid for children < 5 years and 250 mg po bid for children > 5 yr, is given to children with sickle cell disease or hypogammaglobulinemia or after splenectomy to prevent overwhelming pneumococcal infections. Although prophylaxis is controversial, some authorities recommend it until 5 yr of age or 6 yr after surgical removal of the spleen, whichever is longer. The 250-mg bid oral doses of penicillin V and amoxicillin have also been used in adults after splenectomy, but their efficacy is undetermined. To reduce infection rates in patients with chronic granulomatous disease, TMP-SMX in an oral dose of 8 mg/kg (TMP) and 40 mg/kg (SMX) bid for children and 80 mg/kg (TMP) and 400 mg/kg (SMX) for adults is widely used. Specific prophylactic guidelines exist for persons with AIDS and other causes of immunodeficiency.

### Prophylaxis Against Endogenous Pathogens

Prevention of infection by endogenous pathogens, transient or permanent constituents of the normal human flora, usually involves prolonged treatment of chronically colonized persons to prevent spread of bacteria from the site of colonization to more vulnerable sites. For example, adult patients with recurrent skin infections caused by *S. aureus* have fewer infections while receiving orally administered clindamycin in a dose of

150 mg once daily. Hemodialysis patients with nasal carriage of *S. aureus* and a history of recurrent infections caused by this organism have reduced rates of infection orally administered rifampin in a dose of 600 mg bid for 5 days plus topical bacitracin ointment on the nares for 7 days every 3 mo if nasal cultures are positive for *S. aureus*. Mupirocin ointment may substitute for bacitracin in this regimen.

Children with recurrent otitis media experience fewer recurrences when treated prophylactically. Likewise, women with more than three UTIs per year have lower infection rates when treated prophylactically. Patients with compromised host defenses and frequent recurrences of infection appear to have reduced rates of infection when taking TMP-SMX or ciprofloxacin 500 mg bid po.

In other cases, only a day or two of antimicrobial therapy is required to prevent infection. *Streptococcus agalactiae* infection in pregnant women and newborns is reduced with intrapartum penicillin and ampicillin therapy. Generally, only small doses of antimicrobial drugs are recommended to prevent endocarditis in patients with valvular or congenital heart disease when undergoing dental, surgical, or other procedures that are likely to induce bacteremia (see Ch. 208).

Patients undergoing intensive chemotherapy for malignant disease who have absolute neutrophil counts < 500 cells/ $\mu$ L are at high risk for bacterial infection caused by resident flora of the GI tract. Although controversial, prophylactic antimicrobial therapy has been used to suppress the endogenous flora of the bowel to reduce the rate of infection during the neutropenic phase. Antimicrobial resistance patterns and trends in some medical centers argue against this approach to prophylaxis. When used, prophylactic therapy for neutropenic patients includes either oral nystatin (200,000-U lozenge qid or 500,000-U lozenge tid) or clotrimazole (10 mg tablet qid) or clotrimazole (10 mg tablet 6 times/day) to combat *Candida* and other yeasts plus oral antimicrobials. Oral penicillin V 500-mg bid is taken with either SMX/TMP 1600/320 mg bid or ciprofloxacin 500 mg bid or ofloxacin 400 mg bid. Adding fluconazole to either regimen is beneficial.

5

150 mg once daily. Hemodialysis patients with nasal carriage of *S. aureus* and a history of recurrent infections caused by this organism have reduced rates of infection when orally administered rifampin in a dose of 150 mg bid for 5 days plus topical bacitracin ointment for 7 days every 3 mo if nasal cultures are positive for *S. aureus*. Mupirocin ointment applied intranasally may substitute for bacitracin in this regimen.

Children with recurrent otitis media experience fewer recurrences when treated prophylactically. Likewise, women with more than three UTIs per year have reduced infection rates when treated prophylactically. Patients with compromised intestinal tracts and frequent recurrences of ascending infection appear to have reduced rates of infection when taking TMP-SMX or ciprofloxacin 500 mg bid po.

In other cases, only a day or two of antimicrobial therapy is required to prevent infection. *Streptococcus agalactiae* infection in pregnant women and newborns can be reduced with intrapartum penicillin or ampicillin therapy. Generally, only low doses of antimicrobial drugs are recommended to prevent endocarditis in patients with valvular or congenital heart disease when undergoing dental, surgical, or other procedures that are likely to induce bacteremia (see Ch. 208).

Patients undergoing intensive chemotherapy for malignant disease who have absolute neutrophil counts < 500 cells/ $\mu$ L for  $\geq 2$  days are at high risk for bacterial infection caused by resident flora of the GI tract. Although controversial, prophylactic antimicrobial therapy has been used to suppress the endogenous flora of the bowel and reduce the rate of infection during the neutropenic phase. Antimicrobial resistance patterns and trends in some medical centers argue against this approach to prophylaxis. When used, prophylactic therapy for neutropenic patients includes either oral nystatin (200,000-U lozenge qid or 500,000-U solution or tablet qid) or clotrimazole (10 mg troche 5 times/day) to combat *Candida* sp. and other yeasts plus oral antimicrobial drugs. Oral penicillin V 500 mg bid is taken with either SMX/TMP 1600/320 mg bid or a fluoroquinolone—ciprofloxacin 500 mg bid, or ofloxacin 400 mg bid, or ofloxacin 400 mg bid. Adding fluconazole to either regimen is not beneficial.

In patients undergoing allogeneic bone marrow transplantation, similar approaches are used to prevent infection during the prolonged phase of neutropenia before marrow engraftment.

Critically ill patients in ICUs who require intubation and mechanical ventilation for more than a few days are at high risk for developing bacterial pneumonia. Selective decontamination of the digestive tract, which is used extensively in Europe, combats this risk with topical and systemic antimicrobial therapy. It is designed to prevent oropharyngeal and GI colonization by aerobic gram-negative bacilli and *Candida* sp. In most studies, a paste containing polymyxin B, tobramycin or gentamicin, and amphotericin B or nystatin has been applied to the oral mucosa qid for the duration of intubation, and solutions of the same antimicrobial drugs have been administered orally or through a nasogastric tube several times a day. Cefotaxime 1 to 2 g IV q8h has also been administered for the first 5 days. Although such regimens have reduced infection rates in several studies, especially those for nosocomial pneumonia, there is no consensus recommending its use in the USA.

## CHAPTER 153 - ANTIBACTERIAL DRUGS / 1125

## Prophylaxis in Surgery

Antimicrobial drugs are used perioperatively to prevent endogenous flora from gaining entry into normally sterile body sites. As a rule, prophylaxis is beneficial in so-called clean procedures only when prosthetic material or devices are being inserted and in clean-contaminated procedures, which are defined by transections of respiratory, GI, or GU tract mucosal surfaces. All of these surfaces are populated by the host's normal flora, which inevitably contaminate the wound and cause a high incidence of wound infections; if prophylactic antimicrobial drugs are not administered.

Selection of antimicrobial drugs is based on the bacterial most likely to contaminate the wound during a specific procedure. Commonly recommended prophylaxes for various procedures are listed in TABLE 153-3. These recommendations may need to be modified in light of patient allergies, local resistance patterns, and local infection rates.

To effectively prevent surgical site infections, antimicrobial drugs are usually administered IV at the induction of anesthesia to ensure adequate concentrations in the wound when the first incision is made. De-

TABLE 153-3. ANTIMICROBIAL PROPHYLAXIS IN SURGERY

Category and Procedure	Adult Dosage*
<b>Neurosurgical</b>	
Craniotomy—high risk only, eg, re-explorations, microsurgery, and entry into sinuses or nasopharynx	Vancomycin 1 g IV and gentamicin 1.5 mg/kg IV preop; or cefazolin 1 g IV preop
CSF shunt placement—only in hospitals with high infection rates (15–20%)	Trimethoprim 160 mg IV and sulfamethoxazole 800 mg IV preop and q 12 h for 3 doses; or vancomycin 10 mg and gentamicin 3 mg injected into cerebral ventricle
<b>Ophthalmic</b>	
Extraction of lens, including insertion of prosthesis	Gentamicin, tobramycin, or neomycin-gramicidin-polymyxin B drops over 2–24 h; and cefazolin 100 mg subconjunctivally at end of procedure
<b>Otolaryngologic</b>	
Major head and neck surgery involving mucosa of oral cavity or pharynx	Cefazolin 1–2 g IV; or clindamycin 600–800 mg IV $\pm$ gentamicin 1.5 mg/kg IV preop and q 8 h for 2 doses
<b>Cardiac</b>	
Median sternotomy, coronary artery bypass grafting, valve surgery, and pacemaker insertion	Cefazolin 2 g or cefuroxime 1.5 g IV preop and q 4–6 h intraoperatively; or vancomycin 1 g preop

Table continues on the following page.



## 1126 / SECTION 13 - INFECTIOUS DISEASES

TABLE 153-3. ANTIMICROBIAL PROPHYLAXIS IN SURGERY (Continued)

Category and Procedure	Adult Dosage*
<b>Noncardiac thoracic</b> Pneumonectomy, lobectomy, and other resections or esophageal operation	Cefazolin 1-2 g IV preop and q 6 h for 24 h vancomycin 1 g IV preop
<b>Abdominal</b> Gastroduodenal—with hemorrhage, malignancy, obstruction, and other high-risk features Gastric bypass Percutaneous gastrostomy Biliary tract (including ERCP)—with acute symptoms, previous surgery, jaundice, and other high risk features Appendectomy	Cefazolin 1-2 g IV preop or clindamycin 600 mg IV preop and gentamicin 120 mg IV preop Cefazolin 1-2 g IV preop Cefazolin 1-2 g IV preop Cefazolin 1-2 g IV preop; or gentamicin 120 mg IV preop and q 8 h for 3 doses Cefoxitin or cefotetan or cefmetazole 1 g IV preop and q 6 h for 3 doses if not penicillin or metronidazole 500 mg and gentamicin 120 mg/kg preop
<b>Colorectal</b>	Neomycin 1 g and erythromycin base 1 g PO 2, and 11 PM on day before surgery if not parenteral agent(s) listed below; if enoxacin or cefotetan or cefmetazole IV and q 4 h for 3 doses; or metronidazole 500 mg IV and gentamicin 1.7 mg/kg IV preop and for 3 doses
<b>Obstetric-gynecologic</b> Cesarian section—high risk only, eg, premature rupture of membranes Abortion—2nd trimester instillation Abortion—1st trimester, with history of pelvic inflammatory disease, gonorrhea, or multiple partners Hysterectomy, vaginal or abdominal	Cefazolin 1 g IV after clamping cord and 6 h later Cefazolin 1 g IV preop and q 6 h for 2 doses Penicillin G 1-2 million U IV preop and 3 h later or doxycycline 100 mg po before procedure and 200 mg 1/2 h afterward Cefazolin 1 g IV preop and q 6 h for 2 doses or doxycycline 200 mg IV preop
<b>Orthopedic</b> Arthroplasty, including replacements Open reduction of fractures Lower limb amputation	Cefazolin 1-2 g IV preop and q 6 h for 3 doses or vancomycin 1 g IV preop Cefazolin 1 g IV preop and q 6 h for 3 doses Cefoxitin 2 g IV preop and q 6 h for 4 doses
<b>Vascular</b> Lower extremity or abdominal arterial surgery; lower extremity amputation for ischemia	Cefazolin 1-2 g IV preop and q 6 h for 24 h or vancomycin 1 g IV preop and 12 h after procedure
<b>Urologic</b> Prostatectomy—if bacteriuria is present Penile prosthesis insertion	Cefazolin 1 g IV preop or other agent selected on basis of susceptibility tests Cefazolin 1 g IV preop

\*Agents, dosages, routes, and frequencies given are representative of current expert recommendations. Cefazolin remains highly favored because of its spectrum of bactericidal activity, long half-life, low toxicity. Alternatives are primarily for patients with beta-lactam allergies.

Adapted from Kernodle DS, Kaiser AB: "Postoperative infections and antimicrobial prophylaxis," in *Principles and Practice of Infectious Diseases*, ed. 4, edited by GL Mandell, JE Bennett, and R Dolin. New York: Churchill Livingstone, 1995, pp. 2742-2756 and from "Antimicrobial prophylaxis in surgery," *The Medical Letter* 39:97-102, 1997.

# **PHYLAXIS IN SURGERY (Continued)**

## **Adult Dosage\***

zolin 1-2 g IV preop and q 6 h for 24 h;  
ncomycin 1 g IV preop

zolin 1-2 g IV preop or clindamycin 600 mg  
d gentamicin 120 mg IV preop

zolin 1-2 g IV preop  
zolin 1-2 g IV preop  
zolin 1-2 g IV preop; or gentamicin 80 mg  
op and q 8 h for 3 doses

ctin or cefotetan or cefmetazole 1-2 g IV  
op and q 6 h for 3 doses if not perforated  
metronidazole 500 mg and gentamicin 1.7  
/kg preop

ycin 1 g and erythromycin base 1 g po  
nd 11 pt on day before surgery if elect  
enteral agent(s) listed below; if emergent  
xin or cefotetan or cefmetazole IV preop  
q 4 h for 3 doses; or metronidazole 500 mg  
nd gentamicin 1.7 mg/kg IV preop and  
3 doses

lin 1 g IV after clamping cord and 6 h and  
later

lin 1 g IV preop and q 6 h for 2 doses  
lin G 1-2 million U IV preop and 3 h later  
xycline 100 mg po before procedure and  
ng 1/2 h afterward

in 1 g IV preop and q 6 h for 2 doses; or  
cycline 200 mg IV preop

in 1-2 g IV preop and q 6 h for 3 doses  
ncomycin 1 g IV preop

in 1 g IV preop and q 6 h for 3 doses  
in 2 g IV preop and q 6 h for 4 doses

n 1-2 g IV preop and q 6 h for 24 h; or  
ncomycin 1 g IV preop and 12 h after preop

n 1 g IV preop or other agent selected on  
of susceptibility tests  
n 1 g IV preop

representative of current expert recommendations:  
bactericidal activity, long half-life, low cost, and  
lactam allergies

fections and antimicrobial prophylaxis," in  
JL Mandell, JE Bennett, and R Dolin. New York  
robial prophylaxis in surgery," *The Medical Letter*

pending on the duration of the procedure  
and the pharmacokinetics of the drugs, ad-  
ditional doses may be required intraopera-  
tively. The need for additional doses after the  
wound has been closed is highly controver-  
sial but is recommended by many authorities  
(see TABLE 153-3). The prophylaxis should  
not be continued > 24 h unless an active  
infection was discovered during surgery.

## **Prophylaxis Against Dormant or Latent Pathogens**

Prevention of infection by dormant or la-  
tent pathogens, which are microorganisms  
already residing in the human host but not  
currently causing disease, requires either  
microbial eradication before immunologic  
defenses wane and allow the organisms to  
proliferate, or ongoing suppression of the  
remaining organisms, thereby precluding  
their multiplication and dissemination. For  
example, prophylaxis with acyclovir dra-  
matically reduces the rate and severity of

## **CHAPTER 154 - ANTIVIRAL DRUGS / 1127**

recurrences in persons with genital herpes  
who are subject to frequent attacks (see Ch.  
164). Similarly, prophylactic therapy with  
isoniazid greatly reduces the likelihood of  
symptomatic disease in persons with latent  
TB infection (see Ch. 157).

Antimicrobial therapy is frequently used  
to prevent infections by latent or dormant  
pathogens in immunocompromised patients.  
Prophylaxis with TMP-SMX prevents *Pneu-  
mocystis carinii* pneumonia in persons  
undergoing intensive chemotherapy for  
various types of malignant disease or trans-  
plantation procedures. TMP-SMX, inhaled  
pentamidine, dapsone, and other drugs are  
used to prevent *P. carinii* pneumonia in pa-  
tients with AIDS (see Ch. 163). Also, patients  
with AIDS, depending on their CD4<sup>+</sup> lym-  
phocyte counts and exposures, often receive  
prophylaxis to prevent symptomatic disease  
caused by opportunistic infections (see Ch.  
163). Similar targets for prophylaxis are en-  
countered in transfusion and solid organ  
transplantation procedures (see Ch. 149).

## **154 / ANTIVIRAL DRUGS**

The development of drugs against HIV has  
resulted in a dramatic expansion of new an-  
tiviral chemotherapeutic drugs. The appli-  
cations of some of these drugs are being eval-  
uated for other viral infections such as  
hepatitis B virus (HBV). Other advances in-  
clude development of drugs with improved  
bioavailability for common infections such  
as herpes simplex virus (HSV) and varicella-  
zoster virus (VZV).

Chemotherapeutic intervention can occur  
before or at the time of viral particle attach-  
ment to host cell membranes, during un-  
coating of viral nucleic acids, by inhibiting a  
cellular receptor or factor required for viral  
replication, or by blocking specific virus-  
coded enzymes and proteins produced in the  
host cells that are essential for viral replica-  
tion but not for normal host cell metabolism.

### **Idoxuridine**

Idoxuridine (IDU) acts by irreversibly re-  
placing thymidine in newly synthesized DNA

and producing an abnormal and essentially  
nonfunctional DNA molecule. The drug acts  
on viral and host cell DNA and is highly toxic  
to host cells. Because of its high systemic  
toxicity, IDU has been limited to topical ther-  
apy of herpes simplex keratoconjunctivitis.  
Two topical ophthalmic preparations are  
available. One drop of a 0.1% solution is in-  
stilled conjunctivally q 1 h during waking  
hours and q 2 h at night. Treatment should  
be continued for 5 to 7 days after complete  
healing to lessen the chance of recurrence.  
IDU may cause irritation, pain, pruritus, and  
inflammation or edema of the eyelids; rare  
allergic reactions and photophobia have also  
been reported.

### **Vidarabine**

Vidarabine (adenine arabinoside, ara-A)  
interferes with viral DNA synthesis and is  
effective in the treatment of HSV infections.  
Vidarabine appears less susceptible to the  
development of drug-resistant viral strains



**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**